arXiv:1412.5274v1 [math.CA] 17 Dec 2014

A NOTE ON HARDY q-INEQUALITIES

PENG GAO

Abstract. We give a simple proof of a recently result concerning Hardy q-inequalities.

1. Introduction

For $p \geq 1$, let l^p be the Banach space of all complex sequences $\mathbf{x} = (x_n)_{n\geq 1}$ with norm

$$\|\mathbf{x}\|_p := (\sum_{n=1}^{\infty} |x_n|^p)^{1/p} < \infty.$$

Let $C = (c_{n,k})$ be a matrix acting on the l^p space, the l^p operator norm of C is defined as

$$\|C\|_{p,p} = \sup_{\|\mathbf{x}\|_p=1} \Big|\Big|C\mathbf{x}\Big|\Big|_p.$$

The celebrated Hardy's inequality ([2, Theorem 326]) asserts that for $p > 1$,

$$\sum_{n=1}^{\infty} \Big|\frac{1}{n}\sum_{k=1}^{n} x_k\Big|^p \leq \Big(\frac{p}{p-1}\Big)^p \sum_{n=1}^{\infty} |x_n|^p.$$

Let $0 < q < 1$ be fixed. The definite q-integral or the q-Jackson integral (see[3]) of a function $f : [0, b) \to R, 0 < b < \infty$, is defined as follows:

$$\int_0^x f(t)dt = (1-q)x\sum_{k=0}^{\infty} q^k f(q^k x), \quad x \in [0, b).$$

In the theory of q-analysis the q-analogue $[\alpha]_q$ of a number $\alpha \in R$ is defined by

$$[\alpha]_q = \frac{1-q^\alpha}{1-q}.$$

Recently, Maligranda, Oinarov and Persson [3] considered the q-analogues of the classical Hardy type inequalities, one of their main results is stated as follows:

Theorem 1.1 ([3, Theorem 2.3]). *Let $0 < q < 1$, $\alpha < 1 - 1/p$. If either $1 \leq p < \infty$ and $f \geq 0$ or $p < 0$ and $f > 0$, then the following inequality*

$$\int_0^1 x^{p(\alpha-1)}\Big(\int_0^x t^{-\alpha} f(t)d_q t\Big)^p d_q x < \frac{1}{[1-1/p-\alpha]_q^p}\int_0^1 f^p(t)d_q t$$

holds and the constant $[1-1/p-\alpha]_q$ is best possible.

2000 *Mathematics Subject Classification.* Primary 26D15.

Key words and phrases. Hardy's inequality.

The author is supported in part by NSFC grant 11371043 and the Fundamental Research Funds for the Central Universities.

Using the above result, it is derived in [3, (27)] (here we replace $q^{1-1/p-\alpha}$ in [3, (27)] by q) that if $0 < q < 1$ and either $p > 1$ or $p < 0$, then for $a_n > 0$,

$$\sum_{n=1}^{\infty} \Big(\frac{1}{q^n} \sum_{k=n}^{\infty} q^k a_k\Big)^p \leq \frac{1}{(1-q)^p} \sum_{n=1}^{\infty} a_n^p, \tag{1.1}$$

where the constant $(1-q)^{-p}$ is best possible.

In fact, it is easy to see that Theorem 1.2 is equivalent to inequality (1.1). It is therefore our goal in this note to give a simple proof of Theorem 1.2 by proving inequality (1.1). We only consider the case $p > 1$ here and we prove in the next section the following generalization of inequality (1.1):

Theorem 1.2. *Let $p > 1$, let $\{a_n\}_{n=1}^{\infty}$ be a sequence of non-negative numbers with $a_1 > 0$ and $S = \sum_{n=1}^{\infty} a_n < \infty$. Let $A = (a_{i,j})$ be an upper triangular matrix satisfying $a_{i,j} = a_{j-i+1}, j \geq i$. Then $\|A\|_{p,p} = S$.*

2. Proof of Theorem 1.2

We first note the following generalization of the well-known Schur's test:

Lemma 2.1 ([1, Theorem 1])**.** *Let $p > 1$ be fixed and let $A = (a_{i,j})$ be a matrix with non-negative entries. If there exist positive numbers U_1, U_2 and a positive matrix $(b_{i,j})$, such that*

$$\begin{aligned} \sum_j a_{i,j} b_{i,j}^{1/p} &\leq U_1, \\ \sum_i a_{i,j} b_{i,j}^{-1/q} &\leq U_2. \end{aligned}$$

Then

$$\|A\|_{p,p} \leq U_1^{1/q} U_2^{1/p}.$$

We now apply the above lemma to the matrix A given in Theorem 1.2 by taking $b_{i,j} = 1$ for all i, j. Noting that in this case $U_1 = S, U_2 \leq S$, we obtain immediately $\|A\|_{p,p} \leq S$.

To see that the constant S is best possible, we note that for any given $\epsilon > 0$, there exists an integer $N > 0$ such that $\sum_{i=1}^{N} a_i > S - \epsilon$. Now consider the sequence $\mathbf{x}$ with $x_i = 1, 1 \leq i \leq M$ for some integer $M > 1$ and $x_i = 0, i > M$. When $M > N$, the first $M - N + 1$ rows of $A\mathbf{x}$ have row sum $\geq \sum_{i=1}^{N} a_i > S - \epsilon$. It follows that $\|A\mathbf{x}\|_p / \|\mathbf{x}\|_p > S - 2\epsilon$ when M is large enough. As ϵ is arbitrary, this shows that the constant S is best possible and the proof of Theorem 1.2 is completed.

References

[1] D. Borwein and A. Jakimovski, Matrix operators on l^p, *Rocky Mountain J. Math.*, **9** (1979), 463–477.
[2] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge Univ. Press, 1934.
[3] L. Maligranda, R. Oinarov and L.-E. Persson, On Hardy q-inequalities, arXiv:1403.6292.

Department of Mathematics, School of Mathematics and System Sciences, Beijing University of Aeronautics and Astronautics, P. R. China
E-mail address: penggao@buaa.edu.cn